

Anaerobic degradation of crude oil by sulphate reducing bacteria isolated from soils contaminated with petroleum hydrocarbons

Ahmad Abd Burghal Al- Asadi

Basrah university - College of Science - Biology department

Abstract

This study included isolation a mix cultures of sulphate reducing bacteria from soils in Shaeba regain contaminated with petroleum hydrocarbons by using API medium enriched with oxygen reducing agents and saturated with gas (90% N₂) and (10% CO₂) by using sodium lactate as sole carbon source. The result showed that the number of bacteria ranged (4 x 10⁵ – 4.5 x 10⁵) cells / g of soil. Also using crude oil as a sole of carbon source to study the ability of sulphate reducing bacteria on degradation of crude oil under anaerobic condition . The result showed these bacteria have high ability for degradation of crude oil. The percent rate of degradation in culture was (84.4 %) after (54) days incubation and the rate concentration of total petroleum hydrocarbons (TPH) extraction from culture value (22.06) µg / L , while in control was (190.84) µg / L.

Introduction

The bane of industrial progress has been the generation and release into the environment of the huge amounts of toxic compounds which have caused wide spread contamination of the land and water. These chemical have been released into the environment creating countless number of contaminated sites, most widely distributed environmental pollution can be attributed to hydrocarbons contamination, caused by oil tanker accidents, storage tank rupture, transport accidents , and old petrol station (Jain *et al.*, 2005 and Aoshima *et al.*, 2006).

Crude oil is a complex mixture of hydrocarbons , basically composed of aliphatic , aromatic and asphaltene fraction along with nitrogen, sulfur and oxygen contain compounds . (Aoshima *et al.*, 2006). Some of these compounds have been reports to carcinogenic , mutagenic and have immunodulatory effects on humans, animal and plant life (Miller & Maller,1981 and Van-Gesel *et al.* , 2001). Removed of the hydrocarbons from contaminated environments involved physical and chemical processes which expensive and some time difficult to execute, so the microbial

degradation of spilled hydrocarbons is major technique in natural decontamination processes, converting toxic organic to harmless products, often carbon dioxide and water, also its can degrade a wide range of hydrocarbons (Saadoun, 2002 and Koma *et al.*, 2003). The most processes of hydrocarbons biodegradation occurs in aerobic condition but there are several indicating the anaerobic degradation occurs in the absent of oxygen by Sulphate Reducing Bacteria .(Balk, 2007 and Widdel *et al.*, 2007).

The sulphate – reducing bacteria (SRB) are large group of anaerobic organisms that play important role in many biogeochemical processes.(Barton and Hamilton, 2007). The main property of this group is obligate anaerobic bacteria population is their active use of sulphate as a final electron acceptor during anaerobic respiratory and capable of generation hydrogen sulphide (H_2S) from the reduction of sulphate .(Boetius *et al.*,2000 and Sahrani *et al.*, 2008).

SRB utilize very wide spectrum of different low molecular compounds (Lactate, acetate, propionate, succinate, format, pyruvate, ethanol and aliphatic acid) as carbon and energy sources (Caumette, 1993). Sulphate and organic matter concentration, temperature and salinity are the main environmental factors controlling the number and distribution of SRB and the rate of bacterial sulphate reduction (Mudryk *et al.*, 2000). They are widely distributed in nature from anaerobic mud found at the bottom of the ocean to the intestines of humans.

(Gad and White, 1996). Also were found to grow environmental contaminants such as petroleum hydrocarbon constituents (benzene, toluene, ethylbenzene, xylenes and

alkanes), petroleum reservoirs and oil production facilities (Zhang and Young, 1997 and Barton and Hamilton, 2007). SRB using to purify industrial wastewater from heavy metals and in recent years studies have also been made on the use of SRB for the biodegradation of organic matter (Sanir *et al.*, 2001; Kleikemper *et al.*, 2002 and Rezscycka *et al.*, 2004). Hence, the aim of this study was isolated and numerated of SRB from contaminated soil and study their ability for anaerobic degradation of crude oil.

Material and methods

Samples collection

Soil samples were collected from Shaeba reign near the south oil refinery company in Basrah which is highly contaminated with crude oil, the samples collected from depth soil at (15- 30) cm under surface by clean handle, especially in sterile plastic containers and sealed to avoid oxygen contamination and under semi anaerobic condition and transferred immediately to the laboratory (Rooney- Varga *et al.*, 1997).

Isolation and cultivation of sulfate reducing bacteria

SRB were isolated from soil samples by using liquid API medium API, (1975) which is used as selective growth medium, with the following compositions : yeast extract (1 g); $MgSO_4 \cdot 7H_2O$ (0.2 g); $Fe(NH_4)_2(SO_4)_2 \cdot 6H_2O$ (0.2 g); NaCl (1 g); K_2HPO_4 (0.01 g); Ascorbic acid (0.1 g) and Sodium lactate (2.24 g). All of the dry chemical were measured out first and prepared by adding to each a liter of distilled water, the pH was adjusted to (7.2) using (1 M) NaOH solution, medium sterilized by autoclave

under (121°C) and pressure (15) bound / inch² for 15 minute , left to cool at room temperature. This medium treated with added oxygen reducing agents has the following: Sodium dithionate (0.3 g / l) and L – cystein (0.28 g / l) these compounds were sterilized by heat , and saturated under gas phase (90% N₂) with (10% CO₂) before being inoculated with the samples (Rabus *et al.*, 1996 and Teske *et al.* , 1996).

(Figure1).The soiled medium was prepared by added (1.5 % wt / vol) agar. To isolate of SRB from samples (1 g) of sample mixed with (9 ml) of liquid API medium and shake well until it become homogenized, (1ml) of homogenized mat surface layer was transfer to screw tubes containing approximately full of liquid API medium under condition mentioned in isolation, tubes were sealed by screw cover and coated with paraffin tape to prevent diffusion of O₂ into medium, and incubated at (35 °C) until blackening of the medium was recorded as positive for SRB presence. (Hirnes *et al.* , 1999 and Carignan *et al.*, 1994)

Enumerated of SRB

For determination of cell numbers, dilution series (10⁻² - 10⁻⁹) was prepared from each samples with liquid API medium under condition identical to that mention in isolation of SRB in screw stopped tubes. From (10⁻⁵-10⁻⁷) dilutions, (0.5) ml was incubated into triplicate anaerobic roll tubes containing approximately full of API agar medium at (45) °C , tubes were stopped well and coated with paraffin tape , then tubes were rolling between two hand to speared inoculums through the medium, incubated at (35) °C until well formed colonies become visible (Hines *et al.*,

1999), numbers of cells measured as following

Numbers of cells / g of soil = Numbers of colonies × 1/dilution.

Purification of SRB

The colonies obtained from cultivation series dilution in API agar were picked by means of finely drawn sterile Pasteur pipettes , the colonies were immediately transferred into tube of fresh liquid medium , community of SRB formed by mixing equal volumes of eight cultures of bacterial communities originating from various samples (Rabus *et al.*, 1996).

Biodegradation of crude oil by SRB

Using north rumella crude oil in the chemical experiment to determined ability of SRB for degraded crude oil, stored in cool place by using sterilized container until being used. conical flasks (100) ml with (80) ml of sterilized liquid API medium without sodium lactate, added (3%) of crude oil by sterilized pipettes to every conical, then inoculums' by (3) ml of activated mix cultures of SRB , incubated at (35) °C for (54) days in dark without shaking at form semi sloping to maximum the contact area between oil and medium, conical flasks were sealed and coated with paraffin tape , control sample was prepare by added (3%) crude oil to medium only and incubate under same conditions (Rueter *et al.*,1994 and Rabus *et al.*,1996).

Extraction of oil hydrocarbons

The oil consumption ratio by SRB in liquid medium and control samples , estimated by the weight measurement method ,extraction (80) ml growth culture with (100) ml of carbon tetra chloride in separation funnel ,was well

mixing many times, the mixture of solvent and culture was leave to separate to form two layers , the lower layer (contain oil hydrocarbons) was collected and transferred into separation Colum contain an anhydrous sodium sulfate which make up absorbed water and other contaminates , hydrocarbons fraction was collected in round bottle , and

volatilized of solvent by rotary evaporator under low pressure at(55) °C no increase than (60)°C . The weight of oil extraction was measured after drying from solvent (UNEP, 1992) , oil consumption ratio was calculated from residual oil components as the following:

$$\text{Degradation rate}\% = \frac{\text{mg of crude oil control} - \text{mg of crude oil test}}{\text{mg of crude oil control}} \times 100$$

Total Petroleum Hydrocarbons (TPH) was determined in testing and control samples, dissolved dry hydrocarbon extraction with n- hexane (5) ml , then were measured by spectrofluometer system (type Shimadzu- RF540) was equipped with recording (type Shimadzu - DR3), emission at (360) nm and Excitation at (310) nm .The control sample was measured under same condition, results compared with standared curve as shown in figure (5) for north rumela crude oil by making serial dilution.

Results

The bacterial cultures showed good growth anaerobically in API medium and on agar medium that supplied with oxygen reducing agents and presence of sodium lactate as carbon source under (90% N₂) with (10% CO₂). Rapid growth was observed on this medium as it only took about three days for the liquid medium and one - two days on agar medium to turn blacking due to sulfide production , the observation of black color might imply the present of SRB (figure 2).Black colonies characteristics as single isolates were examined on API agar appear (figure 3). SRB were enumerated in (1g) of soil

samples in solid API medium with about (4 x 10⁻⁵ - 4.5 x 10⁻⁵) cells / g of soil.

The results showed high capacity of SRB for growth in presence of crude oil as sole carbon source , the initially clear medium become black , because of bacterial growth and production of iron sulphide (FeS) after sulphate reducing and emulsification of oil , camper with control samples without SRB present (figure 4). Degradation ability percentage for SRB was (88%).after incubation (54 days). Mean total petroleum hydrocarbons (TPH) remaining in cultivated culture for SRB was (22.06 µg / L) while the control was (190.84 µg / L) (Table 1).

Discussion

The cultures of SRB by use liquid API medium with sodium lactate as carbon source and oxygen reducing agents under (90% N₂) with (10% CO₂) condition , showed good growth within three days, because most SRB are growth on lactate. The growth was estimated on the basic of the amount of sulphate reduced in the medium to turn blackening, (Hanselmann *et al.*, 1995 and Rzczycka *et al.*, 2004)

The results of SRB numeration was showed large numbers of these bacteria , this result is referred to ability of SRB for growth in environments contaminated with oil hydrocarbons , and utilized of compounds which found in crude oil which contain amount of SO_4^{2-} necessary for growth as final electron acceptor , this is a good agreement with the findings Rabus *et al.*(1996) and Zhang and Young (1997) , also during the degradation of crude oil , low molecular weight organic acids such as acetate , propionate and butyrate which in turn may serve as carbon sources for SRB (Cozzarelli *et al.*, 1994). More over to numerous studies have shown that SRB genera are known to readily degrade a wide range of organic acids and associated with the degradation of the respective carbon sources in many environments (Widdle and Bak, 1992; Hanselmann *et al.* ,1995; Purdy *et al.*, 1997; Sass *et al.*,1998 and Kuever *et al.* , 2001). SRB number in the soil range between (4×10^5 to 4.5×10^5) cell per g wet soil, this number highest to the extent that is predominated in

hydrocarbon contaminated soil , because of optimum environment for these microbial (Machaughton *et al.*, 1999)

In the present study growth of SRB on crude oil as sole sources of carbon showed value have been degraded in which that (88.4%) of its through ability to growth and turned black , compared with control samples, this result is agreed with (Ruerer *et al.*, 1994 and Rabus *et al.*, 1996) , in which SRB utilize aliphatic and aromatic hydrocarbons directly from oil samples as the only source of organic substance under anoxic condition. The ability for growth wide variety of SRB in crude oil due to their contain many sources of reducing equivalent for sulphate reduction and organic carbon for cell synthesis (Kleikemper *et al.*, 2002).

SRB represents a special enzymic system such as pyrophosphatase , ATP sulphurylase , bisulphite reductase and desulphofucidin (Mudryk *et al.*,2000), therefore these enzymes increase their ability to hydrocarbons degradation and toleration to high rate of contaminants (Safinowski *et al.*,2004 and widdle *et al.*, 2007)

Figures and table :




(Figure 1) Saturated media with N_2 and CO_2




(Figure 2) Growth of SRB in API medium



(Figure 4) A: Degradation of crude oil by SRB



(Figure 3) Isolated colonies of SRB in solid API medium



(Figure 5) Standard curve for Rumella crude oil

(Table 1) Degradation percentage and main Total Petroleum Hydrocarbons (TPH)

SAMPLES	DEGRADATION %	MAIN TPH $\mu\text{G/L}$
SRB cultures	84.4	22.06
Control		190.84

Reference

- American Petroleum Institute ,API. (1975).
 Recommened practical for biological
 analysis of subsurface water. Cited by:
 Bell, R.G. and Lim, C. (1981).
 Corrosion of mild stainless steel by four
 tropical *desulfovibrio desulfuricans*
 strain. J. Cand. Microbbiol., 27: 242-
 245.
- Aoshima, H., Hirase, T., Tada, T.,
 Ichimura,N.,Yamaguchi,H.,Taguchi,
 M. and Myoenzono,T. (2006).
 Improvement of heavy oil
 degradation by *Rhodococcus*
erythropolis. J. Environ. Biotecchnol.,
 5 : 107-109.
- Balk,M.(2007). Life in the absence of
 oxygen: Alternative electron acceptor
 for anaerobic microorganisms in a
 petroleum environment. Turk.
 J.Biol.,31: 59-66.
- Barton, L.L. and Hamilton, W.A. (2007).
 Sulphate - reducing bacteria:
 Environmental, and engineered system.
 Cambridge University Press. 558 pp.
- Boetius, A., Ravensschlag, K., Schubert, C.
 J. *et al.* (2000). A marine microbial
 consortium apparently mediating
 anaerobic oxidation of methane.
 Nature, 407: 623-626.
- Carignan, R.S., Fukui, M., Wilkes, H. and
 Widdel, F. (1994). Use of diffusion
 samples in oligotrophic lake
 sediments –effects of free oxygen in
 sampler material. Limnol. Oceanogr.,
 39 : 468-474.
- Caumette P., (1993). Ecology and
 physiology of phototrophic bacteria and
 sulphate-reducing bacteria in marine
 salterns. Experientia., 49:473–481.
- Cozzarelli, I. M., Baedeker, M.J,
 Eganhouse, R. P. and Goerlitz, D.F.
 (1994).The geochemical evolution of
 low-molecular-weight organic acids
 derived from the degradation of
 petroleum contaminants in
 groundwater. Geochim. Cosmochim.
 Acta., 58:863–877.
- Gadd, G.M., White, C. (1996). Mixed
 sulfate-reducing bacterial cultures for
 bioprecipitation of toxic metals:
 factorial and response-surface analysis
 of dilution rate, sulfate and substrate
 concentration.Microbiol.,142:197.Cited
 by
- Hanselmann, K. W., Kaiser, J.P., Wenk,
 M., Schon, R. and Bachofen,R. (1995).
 Growth on methanol and conversion of
 methoxylated aromatic substratesby
Desulfotomaculum orientis in the

- presence and absence of sulfate. *Microbiol. Res.*, 150:387–400.
- Hines, M.E., Evans, R.S., Genthner, B.R., Willis, S.G., Friedman, S., Rooney-Varga, J.N. and Devereux, R. (1999). Molecular phylogenetic and biogeochemical studies of sulfate-reducing bacteria in the Rhizosphere of *Spartina alterniflora*. *Appl. Environ. Microbiol.*, 65 : 2209-2216.
- Jain, R.K., Kapur, M., Lobana, S., Lal, B., Sarma, P.M., Bhattacharya, D. and Thakur, I.S. (2005). Microbial diversity: Application of microorganisms for the biodegradation of xenobiotics. *Current Science.*, 89:101-112.
- Kleikemper, J., Schroth, M.H., Sigler, V., Schmucki, M., Bernasconi, S.M. and Zeyer, J. (2002). Activity and diversity of sulfate-reducing bacteria in a petroleum hydrocarbon-contaminated Aquifer. *Appl. Environ. Microbiol.*, 68: 1516-1523.
- Koma, D., Sakashita, Y., Kubota, K., Fujii, Y., Hasumi, F., Chung S.Y., and Kubo, M. (2003). Degradation of car engine base oil by *Rhodococcus* sp. NDKK 48 and *Gordonia* sp. NDKY 76 A. *Biosc. Biotechnol. Biochem.*, 67: 1590 –1593.
- Kuever, J., M. Konneke, A. Galushko, and Drzyzga, O. (2001). Reclassification of *Desulfobacterium phenolicum* as *Desulfobacula phenolica* comb. nov. and description of strain Sax(T) as *Desulfotignum balticum* gen. nov., sp. nov. *International. J. Syst. Evol. Microbiol.*, 51:171–177.
- Macnaughton, S. J., Stephen, J. R., Venosa, A. D., David, G. A., Chang, Y. J and White, D. C. (1999). Microbial population changes during bioremediation of an experimental oil spill. *Appl. Environ. Microbiol.*, 65:3566–3574.
- Miller, E. C. and Maller, J. A., (1981). Search for the ultimate chemicals carcinogens and their reaction with cellular macromolecules. *Cancer*. 47:2327–2345.
- Mudryk, Z. J., Podgorska, B. and Bolalek, J. (2000). The occurrence and activity of sulphate-reducing bacteria in the bottom sediments of the Gulf Gdansk. *Oceanol.*, 42: 105-107.
- Purdy, K. J., Nedwell, D.B., Embley, T.M. and Takii, S. (1997). Use of 16S rRNA-targeted oligonucleotide probes to investigate the occurrence and selection of sulfate-reducing bacteria in response to nutrient addition to sediment slurry microcosms from a Japanese estuary. *FEMS Microbiol. Ecol.*, 24:221–234.
- Rabus, R., Fukui, M., Wilkes, H., and Widdel, F. (1996). Degradative capacities and 16S rRNA-targeted whole-cell hybridization of sulfate-reducing bacteria in an anaerobic enrichment culture utilizing alkylbenzenes from crude oil. *Appl. Environ. Microbiol.*, 62:3605–3613.
- Rooney-Varga, J.N., Devereux, R., Evans, R.S. and Hines, M.E. (1997). Seasonal changes in the relative abundance of uncultivated sulphate-reducing bacteria in a salt marsh sediment and rhizosphere of *spartina*

- alterniflora*. Appl. Environ. Microbiol., 63 : 3894-3901.
- Rueter, P., Rabus, R., Wilkes, H. (1994). Anaerobic oxidation of hydrocarbons in crude oil by new types of sulphate-reducing bacteria. Nature., 372, 455-458.
- Rzeczycka, M., Suszek, A. and Blaszczyk, m. (2004). Biotransformation of phosphogypsum by sulphate-reducing bacteria in media containing different zinc salts. Environ., 13 : 209-217.
- Saadoun, I. (2002). Isolation and characterization of bacteria from crude petroleum oil contaminated soil and their potential to degrade diesel fuel. J. Basic Microbiol., 42: 420-428.
- Safinowski, M. and Meckenstock, R. U. (2004). Enzymatic reactions in anaerobic 2-methylnaphthalene degradation by the sulphate-reducing enrichment culture N 47. FEMS Microbiol. Lett, 240: 99-104.
- Sanir, R.K., Peyton, B.M., Brown, L.T. (2001). Copper-induced inhibition of growth of *Desulfovibrio desulfuricans* G20: assessment of its toxicity and correlation with those of zinc and lead. Appl. Environ. Microbiol., 67: 4765.
- Sass, H., Berchtold, M., and Branke, J. (1998). Psychrotolerant sulphate-reducing bacteria from an oxic freshwater sediment, description of *Desulfovibrio cuneatus* sp. nov. and *Desulfovibrio litoralis* sp. nov. Syst. Appl. Microbiol., 21: 212-219.
- Sharani, F.K., Ibrahim, Z., Yahya, A. and Aziz, M. (2008). Isolation and identification of marine sulphate-reducing bacteria, *Desulfovibrio* sp. and *Citrobacter freundii* from pasir Gudang, Malaysia. Science., 47 : 365-371.
- Teske, A., Sigalevich, P., Cotton, Y. and Muyzer, G. (1996). Molecular identification of bacteria from coculture by denaturing gradient gel electrophoresis of 16S Ribosomal DNA fragments as a tool for isolation in pure culture. Appl. Environ. Microbiol., 62 : 4210-4215.
- United Nations Environment Program, UNEP (1992). Determination of petroleum hydrocarbons in sediments. Reference methods for marine pollution studies. No. 20, 75 pp.
- Van-Gestel, and C. A. M. (2001). The use of acute and chronic bioassays to determine the ecological risk and bioremediation efficiency of oil-polluted soils. Environ. Toxicol. Chem., 20:1438-1449.
- Widdel, F., and F. Bak. F. (1992). Gram-negative mesophilic sulfate-reducing bacteria. In Balows, A. (ed.), The prokaryotes : a handbook on the biology of bacteria: ecophysiology, isolation, identification, applications, vol. : 4. Springer, New York, pp. 3352-3378
- Widdel, F., Musat, F., Knittel, K., and Glushko, A. (2007). Anaerobic degradation of hydrocarbons with sulphate as electron acceptor. In Barton, L.L. and Hamilton, W.A. (ed). Sulphate-reducing bacteria: Environmental and engineered

system. Cambridge University Press. pp:
265-303.

Zhang, X. and Young,L.Y. (1997).
Carboxylation as an initial reaction

in the anaerobic metabolism of
naphthalene and phenanthrene by
sulfidogenic consortia. Appl. Environ.
Microbiol., 63:4759–4764.

التكسير اللاهوائي للنفط الخام بواسطة الجراثيم المختزلة للكبريت المعزولة من التربة الملوثة باليديروكاربونات النفطية

احمد عبد برغال الأسدي

جامعة البصرة - كلية العلوم - قسم علوم الحياة

الخلاصة

تضمنت الدراسة الحالية عزل مزارع خليطة من البكتريا المختزلة للكبريت من التربة الملوثة بالنفط الخام من منطقة الشعبية جنوب العراق باستخدام وسط API المدعم بالعوامل المختزلة للأوكسجين والمشبع بغازي ($90\% N_2$) و ($10\% CO_2$) ويوجد لآكتات الصوديوم كمصدر وحيد للكربون . وقد أظهرت النتائج أن أعداد البكتريا تراوحت بين ($10^4 \times 4,5 - 10^5 \times 4$) خلية / غم من التربة. كما تم استخدام مزيج النفط الخام كمصدر وحيد للكربون لدراسة قابلية البكتريا المختزلة للكبريت على تكسير النفط الخام تحت ظروف لاهوائية وأظهرت النتائج وجود قابلية عالية للبكتريا على تكسير النفط الخام ، إذ بلغ معدل النسبة المئوية للتكسير ($84,4\%$) في مزارع الجراثيم بعد (54) يوماً من الحضان ، كما بلغ معدل تركيز الهيدروكاربونات النفطية الكلية المستخلص من مزارع الجراثيم ($22,06$) مايكروغرام / لتر أما في عينات السيطرة فقد بلغ ($190,84$) مايكروغرام / لتر .